WEST Search History



DATE: Thursday, June 03, 2004

Hide?	<u>Set</u> Name	Query	<u>Hit</u> Count
DB=PGPB,USPT; PLUR=YES; OP=ADJ			
	L5	pta-3965 or pta-3961	0
	L4	pta-3965 or pta-3961L3	0
	L3	pta-3956 or pta-3958 or pta-4030 or pta-3959 or pta-3826 or pta-3957 or pta-3965pta-3961	1
	L2	mop2-1 or rmr1-1 or rmr2-1 or rmr7-1 or rmr7-2 or mop1-2ems or rmr6-1 or rmr9-1	1
	L1	mop1-2	1

END OF SEARCH HISTORY

Welcome to STN International! Enter x:x

LOGINID:ssspta1649axm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

```
* * * * * *
                    Welcome to STN International
NEWS
                Web Page URLs for STN Seminar Schedule - N. America
NEWS 2
                "Ask CAS" for self-help around the clock
                Source of Registration (SR) information in REGISTRY updated
NEWS 3 JAN 27
                and searchable
                A new search aid, the Company Name Thesaurus, available in
NEWS
     4 JAN 27
                CA/CAplus
NEWS 5 FEB 05 German (DE) application and patent publication number format
                changes
                MEDLINE and LMEDLINE reloaded
NEWS 6 MAR 03
     7 MAR 03 MEDLINE file segment of TOXCENTER reloaded
NEWS
NEWS 8 MAR 03 FRANCEPAT now available on STN
NEWS 9 MAR 29 Pharmaceutical Substances (PS) now available on STN
NEWS 10 MAR 29 WPIFV now available on STN
NEWS 11 MAR 29 New monthly current-awareness alert (SDI) frequency in RAPRA
NEWS 12 APR 26
                PROMT: New display field available
NEWS 13 APR 26 IFIPAT/IFIUDB/IFICDB: New super search and display field
                available
NEWS 14 APR 26 LITALERT now available on STN
       APR 27
                NLDB: New search and display fields available
NEWS 15
NEWS 16 May 10 PROUSDDR now available on STN
                PROUSDDR: One FREE connect hour, per account, in both May
NEWS 17 May 19
                and June 2004
NEWS 18
        May 12
                EXTEND option available in structure searching
NEWS 19
        May 12
                Polymer links for the POLYLINK command completed in REGISTRY
        May 17
NEWS 20
                FRFULL now available on STN
                STN User Update to be held June 7 and June 8 at the SLA 2004
NEWS 21 May 27
                Conference
                New UPM (Update Code Maximum) field for more efficient patent
NEWS 22
        May 27
                SDIs in CAplus
NEWS 23
        May 27
                CAplus super roles and document types searchable in REGISTRY
NEWS 24 May 27 Explore APOLLIT with free connect time in June 2004
NEWS EXPRESS MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT
             MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
             AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004
NEWS HOURS
             STN Operating Hours Plus Help Desk Availability
             General Internet Information
NEWS INTER
NEWS LOGIN
             Welcome Banner and News Items
             Direct Dial and Telecommunication Network Access to STN
NEWS PHONE
             CAS World Wide Web Site (general information)
NEWS WWW
```

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may

result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 15:42:16 ON 03 JUN 2004

=> file agricola caplus biosis
COST IN U.S. DOLLARS

SINCE FILE TOTAL ENTRY SESSION 0.21 0.21

FULL ESTIMATED COST

FILE 'AGRICOLA' ENTERED AT 15:42:28 ON 03 JUN 2004

FILE 'CAPLUS' ENTERED AT 15:42:28 ON 03 JUN 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 15:42:28 ON 03 JUN 2004 COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC.(R)

=> s mop1-2

L1 0 MOP1-2

=> s mop2-1

L2 2 MOP2-1

=> dup rem 12
PROCESSING COMPLETED FOR L2

L3 2 DUP REM L2 (0 DUPLICATES REMOVED)

=> d 1-2 ti

- L3 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
- TI Genes affecting transgene silencing in maize and the development of plant lines with low levels of transgene silencing for breeding
- L3 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
- TI Comparison between phenomenological and microscopic optical potential in nuclear data evaluations

=> d 1-2 ab

- L3 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
- Transgenic silencing is a little understood process by which genes introduced into plants are turned off or silenced. Genetic screens were designed to identify corn mutants with reduced gene silencing activity. Such mutant corn lines include Mop1-1; Mop1-2EMS; Mop2-1, mop3-1; CC2343, rmr1-1; rmr1-2; rmr2-1; rmr6-1; rmr7-1; rmr7-2; rmr8-1; rmr9-1; Mop1-4; Mop1-5; and rmr11-1 and seeds derived therefrom, the plants are useful for corn breeding programs to produce inbred and hybrid seed with reduced gene silencing activity.
- L3 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN

 AB The chi square (x2) values, which represent the
- The chi square $(\chi 2)$ values, which represent the degree of the agreement of the calculated cross sections with their exptl. values, are calculated for 3 kinds of optical potential, which are resp. the phenomenol. optimal optical potential (OOP) for a specific element, the global phenomenol. optical potential given by Becchetti and Grenlees (BGP) for a large amount of target nuclei and the microscopic optical potential based on Skyrme force (MOP). Some 14 natural elements (each containing 1-4 isotopes) are calculated with 12-20 n incident energies, which are in the 0.1-24 MeV energy range for each element, resp. The calculated average total $\chi 2$ values

are $.\text{hivin}.\chi\text{OOP2} = 0.496$, $.\text{hivin}.\chi$ MOP2 = 1 .150, and .hivin. $\chi BGP2 = 1.355$, from which one obtains the ratio of average deviation of calculated nuclear data from their exptl. values .hivin. Δ MOP/.hivin. Δ OOP = 1.52 and .hivin. $\triangle BGP/.hivin.\triangle OOP = 1.65$. The microscopic optical potential based on Skyrme force, which has AN anal. formalism without any free parameters, is very useful in nuclear data calcns. and evaluations. => d so ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN PCT Int. Appl., 173 pp. CODEN: PIXXD2 => d pi ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN PATENT NO. KIND DATE APPLICATION NO. DATE _____ ____ WO 2002029070 A2 20020411 WO 2001-US31285 20011005 WO 2002029070 C2 20030220 WO 2002029070 A3 20030814 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG AU 2001-96657 20011005 A5 AU 2001096657 20020415 US 2001-972805 US 2002157133 A1 20021024 20011005 => d rmr1-1 'RMR1-1' IS NOT A VALID FORMAT FOR FILE 'CAPLUS' The following are valid formats: ABS ---- GI and AB ALL ----- BIB, AB, IND, RE APPS ----- AI, PRAI BIB ----- AN, plus Bibliographic Data and PI table (default) CAN ----- List of CA abstract numbers without answer numbers CBIB ----- AN, plus Compressed Bibliographic Data DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing FAM ----- AN, PI and PRAI in table, plus Patent Family data FBIB ----- AN, BIB, plus Patent FAM IND ----- Indexing data IPC ----- International Patent Classifications MAX ----- ALL, plus Patent FAM, RE PATS ----- PI, SO SAM ----- CC, SX, TI, ST, IT SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers; SCAN must be entered on the same line as the DISPLAY, e.g., D SCAN or DISPLAY SCAN) STD ----- BIB, IPC, and NCL IABS ----- ABS, indented with text labels IALL ----- ALL, indented with text labels

 L_3

SO

L3

PΙ

IBIB ----- BIB, indented with text labels IMAX ----- MAX, indented with text labels ISTD ----- STD, indented with text labels OBIB ----- AN, plus Bibliographic Data (original) OIBIB ----- OBIB, indented with text labels SBIB ----- BIB, no citations SIBIB ----- IBIB, no citations HIT ----- Fields containing hit terms HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT) containing hit terms HITRN ----- HIT RN and its text modification HITSTR ----- HIT RN, its text modification, its CA index name, and its structure diagram HITSEO ----- HIT RN, its text modification, its CA index name, its structure diagram, plus NTE and SEQ fields FHITSTR ---- First HIT RN, its text modification, its CA index name, and its structure diagram FHITSEQ ---- First HIT RN, its text modification, its CA index name, its structure diagram, plus NTE and SEQ fields KWIC ----- Hit term plus 20 words on either side

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

OCC ----- Number of occurrence of hit term and field in which it occurs

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.
ENTER DISPLAY FORMAT (BIB):ti

L3 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
TI Genes affecting transgene silencing in maize and the development of plant
lines with low levels of transgene silencing for breeding

=> d ti

L4 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN
TI Genes affecting transgene silencing in maize and the development of plant
lines with low levels of transgene silencing for breeding

=> d ti

L5 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN
TI Genes affecting transgene silencing in maize and the development of plant
lines with low levels of transgene silencing for breeding

=> s rmr7-1 L6 1 RMR7-1 => d ti ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN L6 Genes affecting transgene silencing in maize and the development of plant TI lines with low levels of transgene silencing for breeding => s mop 2-12 MOP2-1 L7 => s rmr7-2 or rmr 7-22 RMR7-2 OR RMR 7-2 L8=> d 1-2 ti ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN L8 Genes affecting transgene silencing in maize and the development of plant TТ lines with low levels of transgene silencing for breeding ANSWER 2 OF 2 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN L8STRATEGIES FOR SURVIVAL OF SMALL MAMMALS IN A COLD ALPINE ENVIRONMENT II. TТ SEASONAL CHANGES IN THE CAPACITY OF NONSHIVERING THERMOGENESIS IN OCHOTONA-CURZONIAE AND MICROTUS-OECONOMUS. => s mop1-2ems1 MOP1-2EMS Ь9 => d ti ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN Ь9 Genes affecting transgene silencing in maize and the development of plant TIlines with low levels of transgene silencing for breeding => s rmr6-1 1 RMR6-1 L10 => d ti L10 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN Genes affecting transgene silencing in maize and the development of plant lines with low levels of transgene silencing for breeding => s pta-3965 L11 0 PTA-3965 => s pta-3826 0 PTA-3826 L12 => s pta-3959 0 PTA-3959 L13 => s pta-4030 0 PTA-4030 L14 => s pta-3958 L15 0 PTA-3958

=> s pta-3956

L16

0 PTA-3956

=> s ((chandler, v?) or (chandler v?))/au L17 232 ((CHANDLER, V?) OR (CHANDLER V?))/AU

=> s l17 and paramutaton
L18 0 L17 AND PARAMUTATON

=> del 118 y

=> s l17 and paramutation L18 40 L17 AND PARAMUTATION

=> dup rem 118
PROCESSING COMPLETED FOR L18
L19 21 DUP REM L18 (19 DUPLICATES REMOVED)

=> d 1-10 ti

- L19 ANSWER 1 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN
- TI Genes affecting transgene silencing in maize and the development of plant lines with low levels of transgene silencing for breeding
- L19 ANSWER 2 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 1
- TI A mutation that prevents **paramutation** in maize also reverses Mutator transposon methylation and silencing.
- L19 ANSWER 3 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN DUPLICATE 2
- TI Differential chromatin structure within a tandem array 100 kb upstream of the maize b1 locus is associated with **paramutation**
- L19 ANSWER 4 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 3
- TI The regulatory regions required for B' paramutation and expression are located far upstream of the maize b1 transcribed sequences.
- L19 ANSWER 5 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN
- TI Long-distance cis and trans interactions mediate paramutation
- L19 ANSWER 6 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN DUPLICATE 4
- TI Genetic factors required to maintain repression of a paramutagenic maize pll allele
- L19 ANSWER 7 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
- TI Gene activation and gene silencing.
- L19 ANSWER 8 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
- TI Epigenetic control of gene expression in plants.
- L19 ANSWER 9 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 5
- TI mediator of paramutation1 Is required for establishment and maintenance of paramutation at multiple maize loci.
- L19 ANSWER 10 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 6
- TI Paramutation alters regulatory control of the maize pl locus.

- L19 ANSWER 2 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 1
- Both paramutation and Mutator (Mu) transposon inactivation AB involve heritable changes in gene expression without concomitant changes in DNA sequence. The mechanisms by which these shifts in gene activity are achieved are unknown. Here we present evidence that these two phenomena are linked mechanistically. We show that mutation of a gene, modifier of paramutation 1 (mop1), which prevents paramutation at three different loci in maize, can reverse methylation of Mutator elements reliably. In mop1 mutant backgrounds, methylation of nonautonomous Mu elements can be reversed even in the absence of the regulatory MuDR element. Previously silenced MuDR elements are reactivated sporadically after multiple generations of exposure to mop1 mutations. MuDR methylation is separable from MuDR silencing, because removal of methylation does not cause immediate reactivation. The mop1 mutation does not alter the methylation of certain other transposable elements including those just upstream of a paramutable b1 gene. Our results suggest that the mop1 gene acts on a subset of epigenetically regulated sequences in the maize genome and paramutation and Mu element methylation require a common factor, which we hypothesize influences chromatin structure.

=> d 2 so

- L19 ANSWER 2 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 1
- Proceedings of the National Academy of Sciences of the United States of America, Apr 30, 2002. Vol. 99, No. 9. p. 6130-6135
 Publisher: Washington, D.C.: National Academy of Sciences,
 CODEN: PNASA6; ISSN: 0027-8424

=> d 3 ab

L19 ANSWER 3 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN DUPLICATE 2 Recombination mapping defined a 6-kb region, 100 kb upstream of the ABtranscription start site, that is required for B-I enhancer activity and paramutation-a stable, heritable change in transcription caused by allele interactions in maize (Zea mays). In this region, B-I and B' (the only b1 alleles that participate in paramutation) have seven tandem repeats of an 853-bp sequence otherwise unique in the genome; other alleles have one. Examination of recombinant alleles with different nos. of tandem repeats indicates that the repeats are required for both paramutation and enhancer function. The 6-kb region is identical in B-I and B', showing that epigenetic mechanisms mediate the stable silencing associated with paramutation. This is the first endogenous gene for which sequences required for paramutation have been defined and examined for methylation and chromatin structure. tandem repeat sequences are more methylated in B-I (high expressing) relative to B' (low expressing), opposite of the typical correlation. Furthermore, the change in repeat methylation follows establishment of the B' epigenetic state. B-I has a more open chromatin structure in the repeats relative to B'. The nuclease hypersensitivity differences developmentally precede transcription, suggesting that the repeat chromatin structure could be the heritable imprint distinguishing the two transcription states.

=> d 3 so

L19 ANSWER 3 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN DUPLICATE 2 SO Genes & Development (2002), 16(15), 1906-1918 CODEN: GEDEEP; ISSN: 0890-9369

=> d 7 ab

L19 ANSWER 7 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

=> d 7 so

L19 ANSWER 7 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN SO Plant Physiology (Rockville), (January, 2001) Vol. 125, No. 1, pp. 145-148. print.

CODEN: PLPHAY. ISSN: 0032-0889.

=> d 11-21 ti

- L19 ANSWER 11 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 7
- TI Paramutation in maize.
- L19 ANSWER 12 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 8
- TI Paramutation and related allelic interactions.
- L19 ANSWER 13 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN TI B And pl paramutation in maize: Heritable transcription states programmed during development.
- L19 ANSWER 14 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN
 TI b and pl paramutation in maize: heritable transcription states programmed during development
- L19 ANSWER 15 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 9
- TI Sequences required for **paramutation** of the maize b gene map to a region containing the promoter and upstream sequences.
- L19 ANSWER 16 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 10
- TI Allelic interactions heritably alter the activity of a metastable maize pl allele.
- L19 ANSWER 17 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN TI Paramutation: An allelic interaction that causes heritable changes in transcription.
- L19 ANSWER 18 OF 21 CAPLUS COPYRIGHT 2004 ACS on STN
- TI Paramutation in maize and related allelic interactions

- L19 ANSWER 19 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
- TI **Paramutation**: An allelic interaction that causes heritable changes in transcription.
- L19 ANSWER 20 OF 21 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
- TI Paramutation in maize: Allelic interactions associated with heritable changes in transcription.
- L19 ANSWER 21 OF 21 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

 (2004) on STN DUPLICATE 11
- TI Paramutation, an allelic interaction, is associated with a stable and heritable reduction of transcription of the maize b regulatory gene.